

Genetic Assessment of Adults with Intellectual and Developmental Disabilities

Frequently Asked Questions

Introduction

Many adults whose intellectual and developmental disabilities (IDD) are of unknown origin may benefit from etiologic assessment or reassessment. Etiologic assessment may be helpful in planning preventive care, treatment, and management strategies.

This information sheet answers common questions that primary care clinicians may have about including genetic testing professionals in their patients' care. It provides information for finding the nearest genetic clinic, indications for referral and how to make a referral, and outlines information to give the person with an intellectual and developmental disability and their support persons regarding the referral process.

Frequently Asked Questions

How do I find the nearest genetic clinic for my patient?

- ▶ Contact a local genetic clinic for help in deciding whether to refer, including referral criteria and protocol. Contact information for genetic clinics is available at the American College of Medical Genetics and Genomics website: <https://clinics.acmg.net/>.

What is the likelihood of a patient having a genetic etiology for his/her IDD, to prioritize whom I should consider referring?

- The chances of having a genetic etiology for IDD are generally increased if there is:
- ▶ Severe to profound IDD, congenital malformation(s), epilepsy, or complex medical presentation.
 - ▶ Mild to moderate IDD, congenital malformation(s), epilepsy, or complex medical presentation.
 - ▶ Family history of IDD and/or consanguinity.
 - ▶ Dual diagnosis (IDD and co-occurring mental illness). If possible, patients should be seen by a knowledgeable psychiatrist and/or clinical geneticist who can identify those more likely to have a genetic etiology.

Frequently Asked Questions

Why might a genetic assessment be helpful?

Optimal medical management

- ▶ It may be possible to develop a tailored medical and psychosocial management approach to address physical and mental health issues once the etiology is established. For example, people with Down syndrome have an increased probability of developing thyroid disease throughout their lifespan and will benefit from earlier and more regular screening than guidelines for the general population recommend.
- ▶ Identifying a genetic etiology can have health management consequences for family members. For example, in fragile X syndrome, pre-mutation carrier males and females have the potential to develop fragile X-associated tremor/ataxia syndrome and females have an increased risk of premature ovarian insufficiency.

Family reasons

- ▶ The patient and other family members may want information about the cause of the IDD and the risk of recurrence within the family.
- ▶ There can be substantial guilt about having a child with an IDD. Knowing the cause can relieve parental guilt and provide reassurance. As well, with this knowledge, family members may be able to find support by connecting with syndrome-specific organizations.

Is there a special referral form? What information is needed on the referral?

Contact or check the website of your nearest genetic clinic at the American College of Medical Genetics and Genomics website: <https://clinics.acmg.net/>.

Information to include in a referral:

- ▶ Detailed reason(s) for the referral
- ▶ Any previous genetic test results and the date of the test(s)
- ▶ Copies of other pertinent tests (e.g., brain MRI images, echocardiogram report)
- ▶ Name and contact information of the guardian/conservator/substitute decision maker, if there is one, to provide consent for genetic testing

Are there tests I should do, as a primary care clinician prior to sending a patient with IDD to a genetic clinic?

- ▶ Currently some genetic clinics request that fragile X molecular testing and microarray be done before the patient is seen for genetic assessment. Microarray analysis is a much more sensitive test that has replaced the karyotype (chromosome analysis). Contact the genetics clinic closest to you, for further information as to which tests are appropriate to do prior to referral.
- ▶ The reasons for these tests and for a genetics referral should be discussed with the individual, his/her/their family, and/or the guardian/conservator/substitute decision maker in order to obtain appropriate consent.
- ▶ Salient clinical information should be included on the requisition form.
- ▶ If you order a test, and the results are abnormal or a variant found on microarray or pre-mutation carrier status on fragile X testing, consult a clinical geneticist regarding implications and for patient and family feedback and counseling. Referral to a genetics clinic is highly recommended. In some cases, parental studies may be undertaken through the genetics clinic to assist in interpretation of results.

Is genetic testing covered by health insurance?

- ▶ Genetic assessment and genetic tests are generally covered by medical insurance plans, however, checking with the patient's insurance plan would be best.
- ▶ The medical insurance company or diagnostic laboratory may request that prior authorization is obtained for some genetics tests.

Frequently Asked Questions

Which diagnoses can be detected through genetic testing?

- ▶ More than 2,500 genes have been associated with syndromic and non-syndromic IDD, including autism spectrum disorder. This number is continuing to increase as our knowledge expands.
- ▶ Genetic tests can identify single gene pathogenic variants (mutations), whole or partial chromosome duplications and deletions (including microduplications and microdeletions), imprinting defects, and mitochondrial disorders.
- ▶ Relevant to the population with IDD, examples of diagnoses from genetic testing include Down syndrome, fragile X syndrome, Prader-Willi syndrome, Williams syndrome, Smith-Magenis syndrome and 22q11.2 deletion syndrome. Some syndrome-specific reviews are available at GeneReviews, University of Washington: www.genereviews.org as well as the [Health Watch Tables](#).

Which diagnoses are not primarily genetically determined?

- ▶ IDD can be caused by factors other than genetic conditions, including infections, exposure to toxins/teratogens, and perinatal hypoxemia or trauma. Some forms of cerebral palsy and fetal alcohol spectrum disorders are examples.
- ▶ Such a diagnosis does not preclude the person from also having a genetic syndrome.

What happens when the patient and family members go for genetic testing and counseling?

- ▶ The patient and family members will meet with a genetic counselor and/or geneticist who will obtain a detailed patient and family medical history, as well as explain the reason for the genetic assessment.
- ▶ A physical exam is usually done by the geneticist. This may include measurement of salient physical features (e.g., facial ones), and photography of such features (with appropriate consent) for the individual's medical record.
- ▶ Possible genetic diagnoses may be reviewed and appropriate tests to help determine a diagnosis will be discussed. Potential complexity in interpreting test results may be reviewed (such as uncertain or unexpected results). Consent to retrieve records may be requested.
- ▶ Information will be given so that the patient and family can provide informed consent with respect to the proposed genetic testing.
- ▶ Lab tests will usually be done, including blood and/or urine tests. In some circumstances it may be helpful to obtain other tests such as a skin punch biopsy, X-rays, ultrasound, CT or MRI scans. Referral to other specialists may also be recommended.

How are test results communicated?

- ▶ Results are normally reported to the referring physician and communicated to the patient, family and/or guardian/conservator/substitute decision maker by the geneticist. This may include providing a genetic diagnosis and offering further counseling.
- ▶ Genetic changes causing IDD can occur sporadically or may be inherited. Potential consequences of a hereditary disorder, the likelihood of developing it or transmitting it to one's children, and whether there are means to prevent it or lessen any of these effects would be discussed.
- ▶ Reproductive options should also be discussed at the request of the patient and/or family members.

How does management of the patient proceed?

- ▶ Management may be through the geneticist, primary care clinician or other specialist, depending on the condition, the needs of the patient, and available resources.
- ▶ If no etiology is determined, periodic reassessment may be recommended.

Supporting materials

- i. American College of Medical Genetics and Genomics (ACMG)**
The ACMG and its members engage in coordinated efforts to improve patient care, establish standards of care and laboratory policy, and educate its members about advances important to their practice. www.acmg.net/
- ii. Gene Reviews**
University of Washington, Seattle www.genereviews.org
- iii. National Coordinating Center for Regional Genetic Networks**
Improving access to genetic services for underserved populations. nccrcg.org
- iv. National Organization of Rare Diseases (NORD)**
NORD advances practical, meaningful, and enduring change so people with rare diseases can live their fullest and best lives. rarediseases.org
- v. Orphanet**
Provides information on rare diseases and orphan drugs. www.orpha.net

References

- Boycott K, Hartley T, Adam S, Bernier F, Chong K, Fernandez BA, et al. The clinical application of genome-wide sequencing for monogenic diseases in Canada: Position statement of the Canadian college of medical geneticists. *J Med Genet.* 2015;52(7):431-7.
- Lee RW, Poretti A, Cohen JS, Levey E, Gwynn H, Johnston MV, et al. A diagnostic approach for cerebral palsy in the genomic era. *Neuromolecular Med.* 2014;16(4):821-44.

Copyright

This document complements “Primary care of adults with intellectual and developmental disabilities: 2018 Canadian consensus guidelines”, published in *Canadian Family Physician*, Volume 64(4): April 2018, p254-279.

Originally published as: *Genetic Assessment of Adults with Intellectual and Developmental Disabilities: Frequently Asked Questions*. Forster-Gibson, C., Developmental Disabilities Primary Care Program of Surrey Place, Toronto, 2019.

Modified with permission of the copyright owner, Surrey Place, Toronto, Canada. This tool was reviewed and adapted for U.S. use by Vanderbilt Kennedy Center, Toolkit Advisory Committee, including primary care providers, self-advocates, and family caregivers.

All rights reserved. The content of this tool may not be reproduced or stored in a retrieval system, or transmitted in any form or by any means without the prior written permission of the copyright owner. All content © Surrey Place, 2023. Contact the IDD Toolkit at iddtoolkit@vumc.org for permission to adapt information and tools to your local practice setting.